

Meson Spectrum and the Glueball

G. Ganbold^{a,b,1}

a) Bogoliubov Laboratory of Theoretical Physics, JINR, 141980, Dubna, Russia

b) Institute of Physics and Technology, 210651, Ulaanbaatar, Mongolia

Abstract

The formation and spectrum of two-particle bound states are investigated within a simple relativistic quantum field model with the Yukawa-type interaction. Within this approach, relatively weakly interacting quarks and gluons form stable bound states under the analytic confinement. By introducing a minimal set of free parameters (the quark masses, the coupling constant and the confinement 'radius') and by solving the Bethe-Salpeter equation with one-gluon exchange, we satisfactorily explain the experimental data for most known mesons and the glueball.

1 Introduction

The conventional theoretical description of colorless hadrons within the QCD implies that they are bound states of quarks and gluons considered under the *color* confinement. Another realisation of the confinement is developed in [1] by using an assumption that QCD vacuum is realized by the self-dual homogeneous vacuum gluon field that can lead to the quark and gluon confinement. Hereby, propagators of quarks and gluons are described by entire analytic functions in the p^2 -complex plane [2], i.e. the *analytic* confinement takes place, although there exists a prejudice to the analytic confinement conception (e.g., [3]).

In our earlier paper [4] we considered simple scalar-field models to clarify the pure role of the AC in formation of the hadron bound states. In doing so, we have demonstrated just a mathematical sketch of calculations of the mass spectrum for the "mesons" consisting of two "scalar quarks" by omitting important quantum degrees of freedom such as the spin, color and flavor. Nevertheless, these models gave a quite reasonable sight to the underlying physical principles of the hadron formation mechanism and have resulted in qualitative descriptions of the spectra of "mesons", their Regge trajectories and "glueballs" [5].

It seems reasonable to consider below a more realistic extension of our approach by taking into account the color, flavor and spin degrees of freedom for constituent quarks and gluons. In doing so, we estimate the spectra of pseudoscalar and vector mesons as well as the glueball lowest state.

2 Model

Conventionally, considering this problem within QCD one deals with complicated and elaborate calculations, because the confinement is achieved as a result of strong inter-

¹E-mail: ganbold@thsun1.jinr.ru

action – by involving high-order corrections [6]. Then, there arises a problem of correct and effective summation of these contributions.

On the other hand, the use of a QFT method is effective when the coupling is not large. Then, lower orders of a perturbative technique can result in satisfactory accurate estimates (e.g., in QCD) of observables. Particularly, one can effectively use the one-gluon exchange approximation.

We consider a relativistic physics and for the hadronization processes use the Bethe-Salpeter equation, because the binding energy is not negligible and the relativistic corrections are considerable. In doing so, we use a minimal set of free parameters.

The Lagrangian of the Yukawa-type interaction reads:

$$\begin{aligned} \mathcal{L} = & \left(\bar{\Psi}_\alpha^i [S^{-1}]_{\alpha\beta}^{ij} \Psi_\beta^j \right) + \frac{1}{2} \left(\phi_\mu^a [D^{-1}]_{\mu\nu}^{ab} \phi_\nu^b \right) + g \left(\bar{\Psi}_\alpha^i (i\gamma_\mu)^{\alpha\beta} t_{ij}^a \Psi_\beta^j \phi^a \right) \\ & + g\Lambda \left(\phi_\mu^a \phi_\nu^b \left[\partial_\mu \phi_\nu^c - \partial_\nu \phi_\mu^c \right] \right) f^{abc}, \end{aligned} \quad (1)$$

where $\Psi_{\alpha f}^a(x) = \Psi_{\text{spin flavour}}^{\text{color}}(x)$ and $\phi_\mu^a(x) = \phi_{\text{vector}}^{\text{color}}(x)$ are the quark and gluon fields.

We guess that the matrix elements of hadron processes at large distance (in the confinement region) are integrated characteristics of the quark and gluon propagators and the interaction vertices. Therefore, taking into account the correct symmetry features is more important than the tiny details of these functions.

Our aim is to suggest the most plain forms of these propagators which keep the essential properties and result in a qualitative and semi-quantative description of the hadron spectra. We consider the following propagators for quarks and gluons:

$$\begin{aligned} \tilde{S}_{\alpha\beta}^{ij}(\hat{p}) &= \frac{\delta^{ij}}{m^2} \{ i\hat{p} + m [1 + \gamma_5 \omega(m)] \}_{\alpha\beta} \cdot e^{-\frac{p^2 + m^2}{2\Lambda^2}}, \\ D_{ab}^{\mu\nu}(x) &= \delta_{ab} \delta^{\mu\nu} \frac{\Lambda^2}{(4\pi)^2} e^{-x^2 \Lambda^2/4}, \end{aligned} \quad (2)$$

where Λ - the confinement energy scale, m - the quark mass for given flavour and $0 < \omega(m) \leq 1$ is a smoothly decreasing function. These are entire analytic functions and could serve reasonable approximations to the explicit propagators in the hadronization region.

Consider the partition function and take an explicit integration over ϕ -variable:

$$\begin{aligned} Z &= \iint \delta\bar{\Psi} \delta\Psi e^{-\mathcal{L}_F[\bar{\Psi}, \Psi]} \int \delta\phi e^{-\mathcal{L}_B[\phi] - \mathcal{L}_{int}[\bar{\Psi}, \Psi, \phi]} \\ &= \iint \delta\bar{\Psi} \delta\Psi e^{-\mathcal{L}_F[\bar{\Psi}, \Psi] - \mathcal{L}_2[\bar{\Psi}, \Psi] - \mathcal{L}_3[\bar{\Psi}, \Psi] - \mathcal{L}_G + \dots}, \end{aligned} \quad (3)$$

where the two-quark and two-gluon bound states in the one-gluon exchange mode are generated by terms

$$\mathcal{L}_2 = \frac{g^2}{2} \int \delta\phi e^{-\mathcal{L}_B[\phi]} \left(\bar{\Psi} \Gamma \Psi \right) D \left(\bar{\Psi} \Gamma \Psi \right), \quad \mathcal{L}_G = \frac{27g^2}{2} \int \delta\phi e^{-\mathcal{L}_B[\phi]} (\phi \phi D \phi \phi). \quad (4)$$

3 Light and Heavy Mesons

Below we shortly describe important steps of our approach for two-quark bound states. First, we allocate the one-gluon exchange between quark currents and go to the relative

co-ordinates in the centre-of-masses system. Then, make a Fierz transformation to get the colorless bilocal quark currents and introduce an orthonormal basis $\{U_Q(x)\}$ indexed by the quantum numbers $Q = \{n\kappa l m\}$, where l stands for the orbital quantum number. We diagonalize $\mathcal{L}_2$ on colorless quark currents and use a Gaussian representation by introducing auxiliary meson fields $B_{\mathcal{N}}$. Then, we apply the Hadronization Ansatz to identify $B_{\mathcal{N}}(x)$ as meson fields with quantum numbers $\mathcal{N} = \{JQff\}$. The partition function for mesons reads:

$$Z_{\mathcal{N}} = \int DB_{\mathcal{N}} e^{-\frac{1}{2}(B_{\mathcal{N}}[1+g^2\text{Tr}V_{\mathcal{N}}SV_{\mathcal{N}}S]B_{\mathcal{N}})+W_I[B_{\mathcal{N}}]}, \quad (5)$$

where S - the quark propagator and $V_{\mathcal{N}} = \Gamma_J \int dy U_Q(y) \sqrt{D(y)} \exp\{\frac{y}{2} \overleftrightarrow{\partial}\}$ is a vertex function. The interaction between mesons is described by $W_I[B_{\mathcal{N}}] = O[B_{\mathcal{N}}^3]$.

The diagonalization of the quadratic form is equivalent to solution of the Bethe-Salpeter equation (in the ladder approximation) on the orthonormal system $U_{\mathcal{N}}$:

$$g^2 \text{Tr}(V_{\mathcal{N}}SV_{\mathcal{N}'}S) = \iint dxdy U_{JQ}(x) g^2 \Pi_p(x, y) U_{J'Q'}(y) = -\delta^{QQ'} \delta^{JJ'} \lambda_{JQ}(-p^2). \quad (6)$$

Then, the meson mass is defined by equation:

$$1 - \lambda_{\mathcal{N}}(M_{\mathcal{N}}^2) = 0. \quad (7)$$

The polarization kernel $\Pi_p(x, y)$ is real and symmetric that allows one to use a variational technique. For simplicity we consider orbital excitations: $n = 0, l \geq 0$. By solving the variational equation for the meson mass

$$\begin{aligned} 1 = & \frac{\alpha_s \Lambda^4}{3\pi m_1^2 m_2^2} \exp \left\{ \frac{M_l^2(\mu_1^2 + \mu_2^2) - (m_1^2 + m_2^2)}{2\Lambda^4} \right\} \frac{2^l}{(l+1)!} \\ & \cdot \max_{0 < b < 1} \left\{ [b(1-b/2)]^{l+2} (-1)^l \frac{d^l}{dA^l} \frac{1}{A^2} \exp \left(-\frac{M_l^2(\mu_1 - \mu_2)^2}{4A\Lambda^2} \right) \cdot \left[\frac{2\rho_J}{A} \right. \right. \\ & \left. \left. + \frac{M_l^2(\mu_1 - \mu_2)^2}{4\Lambda^2} \left(\frac{2}{A} - \frac{1}{A^2} \right) + \frac{M_l^2\mu_1\mu_2 + m_1m_2(1 + \chi_J \cdot \omega(m_1/\Lambda) \cdot \omega(m_1/\Lambda))}{\Lambda^2} \right] \right\}, \\ A = 1 + 2b, \quad \mu_i = \frac{m_i}{m_1 + m_2}, \quad \rho_J = \left\{ 1, \frac{1}{2} \right\}, \quad \chi_J = \{+1, -1\} \quad \text{for } J = \{P, V\} \end{aligned}$$

we fit the known experimental data and the optimal set of parameters is obtained:

$$\begin{aligned} \alpha_s \equiv \frac{g^2}{4\pi} = 0.324, \quad \Lambda = 668\text{MeV}, \quad m_u = m_d = 179\text{MeV}, \\ m_s = 202\text{MeV}, \quad m_c = 781\text{MeV}, \quad m_b = 4324\text{MeV}, \end{aligned} \quad (8)$$

3.1 Meson Ground States

With these parameters first we estimate the pseudoscalar and vector meson masses for the ground state $l = 0$ in the wide range from light $\pi(138)$ to heavy $\Upsilon(9460)$ represented in Table 1. Note, our parameter α_s is close to a particular value $\alpha_s = 0.34$ known from the τ -lepton decay. In the present paper we consider fixed coupling constant in the range from hundred MeV to $\simeq 10\text{GeV}$, although there exists another point of view to introduce a running α_s dependent on the energy scale [8, 9].

$J^{PC} = 0^{-+}$	M_P	$J^{PC} = 0^{--}$	M_P	$J^{PC} = 1^{--}$	M_V	$J^{PC} = 1^{--}$	M_V
$\pi(140)$	140	$D(1870)$	1929	$\rho(770)$	770	$K^*(892)$	891
$\eta(547)$	547	$D_s(1970)$	2012	$\omega(782)$	782	$D^*(2010)$	2001
$\eta_c(2979)$	3053	$B(5279)$	5356	$\Phi(1019)$	1004	$D_s^*(2112)$	2081
$\eta_b(9300)$	9454	$B_s(5370)$	5398	$J/\Psi(3097)$	3097	$B^*(5325)$	5356
$K(495)$	495	$B_c(6400 \pm 400)$	6154	$\Upsilon(9460)$	9460		

Table 1: Estimated masses (in MeV) for the pseudoscalar and vector mesons.

3.2 Meson Excites States

Compared to the ground states, the orbital excitations $l > 0$ take place in larger distances and should be less sensitive to the short-range details of the quark and gluon propagators. A correct description of the mesonic Regge trajectories can serve an additional testing ground for our model, for the approximated propagators. Our estimates (solid lines) for the ρ -meson and K -meson families excitations plotted versus $J = l + S$ is given in Figure 1 in comparison with the experimental data (dots).

As is expected, our model describes well the Regge trajectories of mesons. The slope of the Regge trajectories is very sensitive on the confinement scale parameter Λ and its optimal value is fixed near $\Lambda = 668$ MeV.




Figure 1: M_J^2 plotted versus the quantum number $J = l + S$ (the Regge trajectories) for the ρ - and K^* -meson families. Solid lines correspond to our estimates and dots depict the experimental data from *Particle Data Group 2004*.

4 Glueball

There may exist the glueballs, bound states of gluons without any constituent quarks. The experimental status of glueballs is not clear [10], although it is evident that the structure of QCD vacuum plays the main role in their formation. We suppose that the lowest-mass glueball may be approximated by the pure "gluon-gluon" bound state with scalar quantum numbers $J^{PC} = 0^{++}$.

Omitting details, the equation for the lowest-state glueball mass reads

$$1 - \tilde{\Pi}(-M_G^2) = 0 \quad \Rightarrow \quad M_G = \Lambda \sqrt{2 \ln \left(\frac{2\pi(2 + \sqrt{3})^2}{27 \alpha_s} \right)}. \quad (9)$$

With obtained values $\{\alpha_s = 0.324, \Lambda = 668 \text{ MeV}\}$ we estimate

$$M_G = 1433 \text{ MeV} . \quad (10)$$

This result may be compared with other predictions of the glueball mass:

$$\begin{aligned} M_{G_S} &= 1580 \pm 200 \text{ MeV} & [Y.A.Simonov, 2000] , \\ M_{G_F} &= 1530 \pm 200 \text{ MeV} & [H.Forkel, 2001] . \end{aligned}$$

Concluding we remark that our simple relativistic field model with an analytic confinement of quarks and gluons gives a quite reasonable sight to the underlying physical principles of the hadron formation mechanism. It allows one to describe qualitatively the meson spectra in the wide range from light $\pi(138)$ to heavy $\Upsilon(9460)$. By using only a minimal set of fundamental parameters (the constituent quark masses m_f , the coupling constant α_s and the confinement scale Λ) we are able to describe the meson ground states, the Regge trajectories of orbital excitations and the low-mass glueball state with sufficient accuracy. Hereby, the coupling constant is relatively small that makes the use of one-gluon exchange modes reliable for the consideration.

The author thanks Ts.Baatar, G.V.Efimov, S.B.Gerasimov, E.Klempt, S.N.Nedelko and W.Oelert for useful discussions and comments.

References

- [1] H. Leutwyler, Phys. Lett., **96B** (1980) 154; Nucl. Phys., **B179** (1981) 129.
- [2] G.V. Efimov and S.N. Nedelko, Phys. Rev., **D51** (1995) 174;
G.V. Efimov, A.C. Kalloniatis and S.N. Nedelko, Phys. Rev., **D59** (1999) 014026.
- [3] A. Ahlig et al., Phys. Rev., **D64** (2001) 014004.
- [4] G.V. Efimov and G. Ganbold, Phys. Rev., **D65** (2002) 054012.
- [5] G. Ganbold, AIP Conf. Proc., **717** (2004) 285; **786** (2005) 127.
- [6] M. Baldicchi and G.M. Prospri, hep-ph/0310213 (2003).
- [7] Ja.V. Burdanov et al., Phys. Rev., **D54** (1996) 4483.
- [8] D.V. Shirkov, Theor. Math. Phys., **132** (2002) 484.
- [9] A.V. Nesterenko, Int. J. M. Phys., **A18** (2003) 5475.
- [10] E. Klempt, hep-ph/0404270 (2004).